



ICD-10-AM/ACHI/ACS Eleventh Edition

Errata 1, June 2019 For implementation 1 July 2019

ICD-10-AM Tabular List

Page 60

Add Code also instruction

D68.3 Haemorrhagic disorder due to circulating anticoagulants

∇ 0303

Haemorrhage during long term use of anticoagulants
Hyperheparinaemia
Increase in:
• anti- VIIIa
• anti- IXa
• anti- Xa
• anti- XIa
• antithrombin

Code also, if applicable:

- [nontraumatic haematoma of skin and subcutaneous tissue \(L98.8\)](#)
- [nontraumatic haematoma of soft tissue \(M79.8-\)](#)

Use additional external cause code (Chapter 20) to identify any administered anticoagulant.

Excludes: abnormal coagulation profile (R79.83)
long term use of anticoagulants without
haemorrhagic disorder (Z92.1)

Page 184

Add Inclusion terms

J17.2* Pneumonia in mycoses
Pneumonia in:
• candidiasis (B37.1†)
• coccidioidomycosis (B38.0–B38.2†)
• histoplasmosis (B39.-†)
• [pneumocystosis \(B48.5†\)](#)

Page 340-1

Add Exclusion term and note

R40
V1905

Somnolence, stupor and coma

Note: For classification purposes, where there are multiple GCS scores documented in an episode of care, code assignment should be based on the lowest GCS score, that is, the most severe level of consciousness impairment.

R40.0

Somnolence

Decreased (level of) consciousness (nontraumatic)
Drowsiness
GCS score 13–15

R40.1

Stupor

GCS score 9–12

Semicoma

Excludes: due to trauma (S06.0-)

stupor:

- catatonic (F20.2)
- depressive (F31–F33)
- dissociative (F44.2)
- manic (F30.2)

that with any head injury classifiable to Chapter 19 (S06.01–S06.05)

R40.2

Coma

GCS score ≤ 8

Loss of consciousness (nontraumatic) NOS

Unconsciousness NOS

Excludes: coma:

- diabetic (E10–E14)
- hepatic (K72.-)
- hypoglycaemic (nondiabetic) (E15)
- neonatal (P91.5)
- that with any head injury classifiable to Chapter 19 (S06.01–S06.05)
- uraemic (N19)
- syncope (R55)

Page 400

Amend Use additional instruction

T78.0 Anaphylaxis and anaphylactic shock due to adverse food reaction

V2115

Use additional external cause code (Y37.0+–Y37.5, Y37.8, Y37.9) to identify allergen, if known.

Page 493

Add ACS number

Z58.7 Exposure to tobacco smoke

[V0050](#), 2118

Passive (involuntary) smoking

Includes: exposure to secondhand tobacco smoke (from):

- cigarette
- pipe
- waterpipe

Excludes: mental and behavioural disorders due to the use of tobacco (F17.-)
newborn affected by maternal use of tobacco (P04.2)
personal history of tobacco use disorder (Z86.43)
tobacco use (Z72.0)

ICD-10-AM Alphabetic Index

Page 113

Add and delete terms, codes and cross reference

Disorder (of) — *see also Disease*

...

- cognitive ([mild](#)) ([moderate](#)) [F06.7](#) ~~NEC R41.8~~

- - due to or secondary to

- - - age (age-associated) R41.8

- - - general medical condition F06.7

~~- - mild (organic) F06.7~~

- - severe — *see Dementia*

- colon K63.9

Page 116

Add terms, code and cross reference

Disorder (of) — *see also Disease*

...

- nervous system G98

- - autonomic (peripheral) G90.9

- - - specified NEC G90.8

- - central G96.9

- - - specified NEC G96.8

- - parasympathetic G90.9

- - specified NEC G98

- - sympathetic G90.9

- - vegetative G90.9

- [neurocognitive \(mild\) \(moderate\) F06.7](#)

- - severe — *see Dementia*

- neurohypophysis NEC E23.3

Page 172

Add and delete terms, codes and cross references

Hepatitis – *continued*

- ...
- viral, virus (without hepatic coma) B19.9
- - with hepatic coma B19.0
- - acute NEC B17.9
- - - specified NEC B17.8
- - - type
- - - - A B15.9
- - - - - with hepatic coma B15.0
- - - - B B16.9
- - - - - with delta-agent coinfection (hepatitis D) (without hepatic coma) B16.1
- - - - - and hepatic coma B16.0
- - - - hepatic coma (without delta-agent coinfection) B16.2
- - - - C B17.1
- - - - D (~~coinfection~~) (acute delta coinfection with chronic hepatitis B ~~with delta-agent~~) (without hepatic coma) ~~B16.1~~ [B17.0](#)
- - - - - with hepatic coma B16.0
- - - - E B17.2
- - chronic NEC B18.9
- - - specified NEC B18.8
- - - type
- - - - B B18.1
- - - - - with delta-agent (hepatitis D) B18.0
- - - - C B18.2
- - - - D ([chronic](#) hepatitis B with delta-agent) B18.0
- - - - E B18.8
- - congenital P35.3
- ...

Page 185

Delete nonessential modifier

- Hypotension** (arterial) (~~chronic~~) (constitutional) I95.9
- chronic I95.8

Page 188

Add terms and code

Impaired, impairment (function)

- ...
- myocardium, myocardial (*see also* *Insufficiency/myocardial*) I50.9
- [neurocognitive NEC R41.8](#)
- rectal sphincter R19.89

ACHI Tabular List

Page 29
Amend code
42705-00 Extraction of crystalline lens with implantation of trans-trabecular drainage device Extraction of crystalline lens with implantation of microstents Note: Performed for glaucoma in conjunction with cataract surgery <u>Code first:</u> • type of cataract extraction (see block [200])

ACHI Alphabetic Index

Page 29
Amend code
Block ... - transversus abdominis plane (TAP) - - for - - - analgesia, postprocedural 92517-01 0 [1912]

Australian Coding Standards (ACS)

Page 11

Amend code title in scenario

0002 ADDITIONAL DIAGNOSIS

...

EXAMPLE 22:

A 55-year-old man presented with lower respiratory infection ~~on background~~ ~~of~~ ~~exacerbating~~ his chronic obstructive pulmonary disease. He was commenced on Bactrim BD and physiotherapy performed. On the second day of the admission, the patient complained of having chest tightness since arrival to the hospital. He described the pain as constant, but not radiating. Nurse consulted the treating clinician over the phone and documented "team doctor advised over the phone to administer PRN GTN 300mcg. ECG was performed. Patient states that pain was not relieved with PRN oral GTN, so was given further GTN 300mcg as per team instruction. Patient remains saturating well and telemetry is in situ". Patient responded well to the treatment and was discharged home.

Principal diagnosis: Chronic obstructive pulmonary disease with acute lower respiratory infection

Additional diagnosis: Chest pain

In this example, the chest pain meets the criteria in ACS 0002 in the episode of care, as clinical consultation was undertaken specifically for the condition (*Note:* Telephone consultation with clear documentation of the information exchange is regarded as clinical consultation).

Page 84

Amend code title in scenario

0234 CONTIGUOUS SITES

...

EXAMPLE 3:

Patient admitted with a mass in the tracheobronchial region. A biopsy was performed via a bronchoscopy under sedation, ASA 2. Histopathology report indicated squamous cell carcinoma (SCC).

Codes: C34.8 *Overlapping lesion of bronchus and lung*

M8070/3 *Squamous cell carcinoma NOS*

41898-04 [544] *Endoscopic [needle] biopsy of bronchus*

92515-29 [1910] ~~General anaesthesia~~ *Sedation*, ASA 29

Amend code title in scenario

0236 NEOPLASM CODING AND SEQUENCING

...

PRIMARY NEOPLASM AS A CURRENT CONDITION

A primary neoplasm is classified as a current condition if the episode of care is for:

- diagnosis or treatment of the primary neoplasm, in any of the following circumstances:
 - initial diagnosis of the primary neoplasm
 - treatment of complications of the primary neoplasm or neoplasm treatment
 - operative intervention to remove the primary neoplasm
 - medical care related to the primary neoplasm, including palliative care (see also ACS 2116 *Palliative care*)
 - recurrence of the primary neoplasm previously eradicated from the same organ or tissue (see also ACS 0237 *Recurrence of malignancy*).
- diagnosis or treatment of a secondary (metastatic) malignancy, regardless of when/if the primary site was previously resected. Assign an additional diagnosis code for the primary neoplasm if known, or C80.- *Malignant neoplasm without specification of site* if the site of the primary neoplasm is unknown or unspecified.
- treatment aimed at stopping progression of the neoplasm, such as:
 - pharmacotherapy or radiotherapy (see also ACS 0044 *Pharmacotherapy* and ACS 0229 *Radiotherapy*)
 - subsequent admissions for wider excision (even if there is no residual neoplasm identified on histopathology)
 - staged surgery for prophylactic removal of a related organ.

~~• treatment of another nonmalignant condition, when the malignancy is a comorbidity that has an affect on the episode of care as per ACS 0002 Additional diagnoses.~~

~~• dental clearance prior to radiotherapy. Assign a code for the condition requiring the procedure as per the criteria in ACS 0001 Principal diagnosis.~~

If the episode of care is for treatment of another nonmalignant condition, the malignancy may be classified as a current condition only if it meets the criteria for code assignment as per ACS 0002 Additional diagnoses.

If the episode of care is for dental clearance prior to radiotherapy, assign a code for the condition requiring the procedure as per the criteria in ACS 0001 Principal diagnosis.

If the episode is for follow-up care, the malignancy may be coded as current or as a past history, dependent on the circumstances surrounding the episode of care. (See also ACS 1204 *Plastic surgery*, ACS 2112 *Personal history* and ACS 2114 *Prophylactic surgery*.)

Where there are multiple metastatic sites, assign a code for each site in order to reflect the severity of the condition.

Delete term in scenario

1511 TERMINATION OF PREGNANCY (ABORTION)

...

EXAMPLE 1:

Patient admitted for suction D&C (GA) for termination of pregnancy at (13/40) ~~weeks~~ due to fetal anencephaly.

Codes: O04.9 *Medical abortion, complete or unspecified, without complication*
 O09.1 *Duration of pregnancy 5–13 completed weeks*
 O35.0 *Maternal care for (suspected) central nervous system malformation in fetus*
 35640-03 [1265] *Suction curettage of uterus*
 92514-99 [1910] *General anaesthesia, ASA 99*

Revise scenario

1521 CONDITIONS AND INJURIES IN PREGNANCY

...

EXAMPLE 6:

A pregnant patient with elevated blood pressure (no diagnosis of hypertension) was admitted by her obstetrician to the obstetric unit for hourly BP (blood pressure) monitoring by midwifery staff. She was [assessed by the dermatologist and then](#) treated with calamine lotion for heat rash during the admission. Her blood pressure returned to normal and her rash was no longer evident, therefore she was discharged home the following day.

Codes:
 O99.8 *Other specified diseases and conditions in pregnancy, childbirth and the puerperium*
 R03.0 *Elevated blood-pressure reading, without diagnosis of hypertension*
 O99.7 *Diseases of the skin and subcutaneous tissue in pregnancy, childbirth and the puerperium*
 L74.0 *Miliaria rubra*